



Eastbourne

Local Tobacco Control Profiles

Introduction

The aim of these profiles is to provide information for local government, health organisations, commissioners and other agencies to monitor the impact of tobacco on local communities, and to monitor the services and initiatives that have been put in place to tackle tobacco related problems.

The indicators contained within the web-tool were selected following consultation with stakeholders and a review of the availability of routine data. They are not exhaustive and should be used alongside local intelligence to gain a full picture of the range of tobacco control initiatives and issues in each locality.

The tobacco control profiles are part of a series of products produced by Public Health England that provide local data alongside national comparisons to support local health improvement.

Contents

Spine Charts

Key Indicators	Page 2
Smoking prevalence in adults	Page 2
Smoking related mortality	Page 3
Smoking related ill health	Page 3

Tartan Rugs

Key Indicators	Page 4
Smoking prevalence in adults	Page 4
Smoking related mortality	Page 5
Smoking related ill health	Page 5

Summary Charts

Key Indicators	Page 7
Smoking prevalence in adults	Page 8
Smoking related mortality	Page 12
Smoking related ill health	Page 13

Trends

Key Indicators	Page 15
Smoking prevalence in adults	Page 16
Smoking related mortality	Page 18
Smoking related ill health	Page 19

www.tobaccoprofiles.info


Produced by Public Health England.

For enquiries please contact tobacco.profiles@phe.gov.uk or call us on 020 765 48167

Spine Charts

How to interpret the spine charts

- Significance compared to England average:
- Significantly worse
 - Significance not tested
 - Not significantly different
 - ◆ Regional average
 - Significantly better



Key Indicators

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking Prevalence in adults - current smokers (APS)	2016	21.3	15.5	25.7		4.9
Smoking Prevalence in adults in routine and manual occupations - current smokers (APS)	2016	44.2	26.5	56.1		0.0
Smoking attributable mortality	2014 - 16		272.0			
Smoking attributable hospital admissions	2015/16		1726			
Supporting information - Deprivation score (IMD 2015)	2015	21.3	21.8	42.0		5.0

Smoking prevalence in adults

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Smoking Prevalence in adults - current smokers (APS)	2016	21.3	15.5	25.7		4.9
Smoking Prevalence in adults - ex smokers (APS)	2016	29.6	26.2	11.4		48.3
Smoking Prevalence in adults - never smoked (APS)	2016	49.1	58.3	42.3		73.3
Smoking Prevalence in adults in routine and manual occupations - current smokers (APS)	2016	44.2	26.5	56.1		0.0
Smoking Prevalence in adults in routine and manual occupations - ex smokers (APS)	2016	23.6	21.9	4.4		45.5
Smoking Prevalence in adults in routine and manual occupations - never smoked (APS)	2016	32.2	51.5	25.0		82.6
Smoking Prevalence in adults - current smokers (IHS)	2014	17.4	18.0	29.8		6.1
Smoking Prevalence in adults - ex-smokers (IHS)	2014	40.8	33.9	53.5		14.4
Smoking Prevalence in adults - never smoked (IHS)	2014	41.8	48.1	32.2		68.3
Smoking prevalence in adults in routine and manual occupations - current smokers (IHS)	2014	28.3	28.0	45.3		6.2
Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS)	2014	48.2	30.8	69.8		11.6
Smoking prevalence in adults in routine and manual occupations - never smoked (IHS)	2014	23.5	41.2	13.5		68.3
Smoking prevalence - current smokers (GPPS)	2015/16	16.0	16.4	24.1		8.6
Smoking prevalence - ex smoker (GPPS)	2015/16	34.3	27.5	14.7		37.4
Smoking prevalence - never smoked (GPPS)	2015/16	49.7	56.1	42.8		70.0
Smoking status at time of delivery (current method)	2016/17	13.5	10.7	28.1		2.3
Smoking status at time of delivery (historical method)	2016/17	13.4	10.5	28.1		2.3


Smoking related mortality

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Deaths from lung cancer	2014 - 16	46.9	57.7	109.8		30.9
Deaths from oral cancer	2014 - 16	5.1	4.6	9.1		2.0
Deaths from chronic obstructive pulmonary disease	2014 - 16	41.7	52.2	102.3		25.3
Potential years of life lost due to smoking related illness	2014 - 16	-	1579			


Smoking related ill health

	Period	Local value	Eng. value	Eng. worst	England Range	Eng. best
Premature births (less than 37 weeks gestation)	2013 - 15	68.4	78.4	107.9		57.1
Low birth weight of term babies	2015	3.5	2.8	4.8		0.8
Cost per capita of smoking attributable hospital admissions	2011/12	42.9	38.0	59.3		23.0
Emergency hospital admissions for COPD	2015/16	217	411	1,089		142
Lung cancer registrations	2013 - 15	69.0	78.9	156.9		39.7
Oral cancer registrations	2013 - 15	14.2	14.5	23.6		6.8


Smoking prevalence in adults continued



Smoking related mortality



Smoking related ill health




Comparison with respect to England value



Note: x - Value Missing

Eastbourne

Smoking related ill health continued




Comparison with respect to England value




Summary Charts

Key Indicators


Smoking Prevalence in adults - current smokers (APS) (2016)




Smoking Prevalence in adults in routine and manual occupations - current smokers (APS) (2016)




Smoking attributable mortality (2014 - 16)



Smoking attributable hospital admissions (2015/16)



Supporting information - Deprivation score (IMD 2015) (2015)




Key
 England value and confidence interval
 Significantly worse
 Not significantly different
 Significantly better
 Significance not tested
↑ Eastbourne
 South East region value and confidence interval


Eastbourne

Smoking prevalence in adults


Smoking Prevalence in adults - current smokers (APS) (2016)




Smoking Prevalence in adults - ex smokers (APS) (2016)




Smoking Prevalence in adults - never smoked (APS) (2016)



Smoking Prevalence in adults in routine and manual occupations - current smokers (APS) (2016)




Smoking Prevalence in adults in routine and manual occupations - ex smokers (APS) (2016)




Key England value and confidence interval Eastbourne South East region value and confidence interval
 Significantly worse Not significantly different Significantly better Significance not tested

Smoking prevalence in adults continued


Smoking Prevalence in adults in routine and manual occupations - never smoked (APS) (2016)




Smoking Prevalence in adults - current smokers (IHS) (2014)




Smoking Prevalence in adults - ex-smokers (IHS) (2014)



Smoking Prevalence in adults - never smoked (IHS) (2014)



Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) (2014)




Key England value and confidence interval Eastbourne South East region value and confidence interval
 Significantly worse Not significantly different Significantly better Significance not tested


Eastbourne

Smoking prevalence in adults continued


Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS) (2014)




Smoking prevalence in adults in routine and manual occupations - never smoked (IHS) (2014)




Smoking prevalence - current smokers (GPPS) (2015/16)



Smoking prevalence - ex smoker (GPPS) (2015/16)




Smoking prevalence - never smoked (GPPS) (2015/16)




Key
 England value and confidence interval
 Eastbourne
 South East region value and confidence interval
 Significantly worse
 Not significantly different
 Significantly better
 Significance not tested

Smoking prevalence in adults continued

Smoking status at time of delivery (current method) (2016/17)



Smoking status at time of delivery (historical method) (2016/17)




Key England value and confidence interval ↑ Eastbourne South East region value and confidence interval
 Significantly worse Not significantly different Significantly better Significance not tested


Eastbourne

Smoking related mortality


Deaths from lung cancer (2014 - 16)




Deaths from oral cancer (2014 - 16)



Deaths from chronic obstructive pulmonary disease (2014 - 16)




Potential years of life lost due to smoking related illness (2014 - 16)




Key England value and confidence interval ↑ Eastbourne South East region value and confidence interval
 Significantly worse Not significantly different Significantly better Significance not tested

Smoking related ill health


Premature births (less than 37 weeks gestation) (2013 - 15)




Low birth weight of term babies (2015)




Cost per capita of smoking attributable hospital admissions (2011/12)



Emergency hospital admissions for COPD (2015/16)



Lung cancer registrations (2013 - 15)




Key England value and confidence interval Eastbourne South East region value and confidence interval
 Significantly worse Not significantly different Significantly better Significance not tested

Eastbourne

Smoking related ill health continued

Oral cancer registrations (2013 - 15)




Key England value and confidence interval ↑ Eastbourne South East region value and confidence interval
Significantly worse Not significantly different Significantly better Significance not tested


Trends

Key Indicators


Smoking Prevalence in adults - current smokers (APS) - %




Smoking Prevalence in adults in routine and manual occupations - current smokers (APS) - %




Smoking attributable mortality - per 100,000



Smoking attributable hospital admissions - per 100,000




Supporting information - Deprivation score (IMD 2015)




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

Smoking prevalence in adults


Smoking Prevalence in adults - current smokers (APS) - %




Smoking Prevalence in adults - ex smokers (APS) - %




Smoking Prevalence in adults - never smoked (APS) - %




Smoking Prevalence in adults in routine and manual occupations - current smokers (APS) - %




Smoking Prevalence in adults in routine and manual occupations - ex smokers (APS) - %




Smoking Prevalence in adults in routine and manual occupations - never smoked (APS) - %




Smoking Prevalence in adults - current smokers (IHS) - %




Smoking Prevalence in adults - ex-smokers (IHS) - %




Smoking Prevalence in adults - never smoked (IHS) - %




Smoking prevalence in adults in routine and manual occupations - current smokers (IHS) - %



Smoking prevalence in adults in routine and manual occupations - ex-smokers (IHS) - %




Smoking prevalence in adults in routine and manual occupations - never smoked (IHS) - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

Smoking prevalence in adults continued


Smoking prevalence - current smokers (GPPS) - %




Smoking prevalence - ex smoker (GPPS) - %




Smoking prevalence - never smoked (GPPS) - %



Smoking status at time of delivery (current method) - %




Smoking status at time of delivery (historical method) - %




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

Smoking related mortality


Deaths from lung cancer - per 100,000




Deaths from oral cancer - per 100,000



Deaths from chronic obstructive pulmonary disease - per 100,000




Potential years of life lost due to smoking related illness - per 100,000




Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested

Smoking related ill health


Premature births (less than 37 weeks gestation) - per 1000




Low birth weight of term babies - %




Cost per capita of smoking attributable hospital admissions - £ per capita




Emergency hospital admissions for COPD - per 100,000



Lung cancer registrations - per 100,000



Oral cancer registrations - per 100,000



Key ● England ● Significantly worse ● Not significantly different ● Significantly better ○ Significance not tested